

ON FIXED POINTS OF PERMUTATIONS

PERSI DIACONIS, JASON FULMAN, AND ROBERT GURALNICK

ABSTRACT. The number of fixed points of a random permutation of $\{1, 2, \dots, n\}$ has a limiting Poisson distribution. We seek a generalization, looking at other actions of the symmetric group. Restricting attention to primitive actions, a complete classification of the limiting distributions is given. For most examples, they are trivial – almost every permutation has no fixed points. For the usual action of the symmetric group on k -sets of $\{1, 2, \dots, n\}$, the limit is a polynomial in independent Poisson variables. This exhausts all cases. We obtain asymptotic estimates in some examples, and give a survey of related results.

This paper is dedicated to the life and work of our colleague Manfred Schocker.

1. INTRODUCTION

One of the oldest theorems in probability theory is the Montmort (1708) limit theorem for the number of fixed points of a random permutation of $\{1, 2, \dots, n\}$. Let S_n be the symmetric group. For an element $w \in S_n$, let $A(w) = \{i : w(i) = i\}$. Montmort [Mo] proved that

$$(1.1) \quad \frac{|\{w : A(w) = j\}|}{n!} \rightarrow \frac{1}{e} \frac{1}{j!}$$

for j fixed as n tends to infinity. The limit theorem (1.1) has had many refinements and variations. See Takács [Ta] for its history, Chapter 4 of Barbour, Holst, Janson [BHJ] or Chatterjee, Diaconis, Meckes [CDM] for modern versions.

The limiting distribution $P_\lambda(j) = e^{-\lambda} \lambda^j / j!$ (In (1) $\lambda = 1$) is the Poisson distribution of “the law of small numbers”. Its occurrence in many other parts of probability (see e.g. Aldous [Al]) suggests that we seek generalizations of (1.1), searching for new limit laws.

In the present paper we look at other finite sets on which S_n acts. It seems natural to restrict to transitive action – otherwise, things break up into orbits

Date: August 15, 2007.

1991 Mathematics Subject Classification. 20B30, 20B35, 05A16, 60C07.

Key words and phrases. Fixed point, derangement, primitive action, O’Nan-Scott theorem.

Diaconis was supported by NSF grant DMS-0505673. Fulman received funding from NSA grant H98230-05-1-0031 and NSF grant DMS-0503901. Guralnick was supported by NSF grant DMS-0653873.

We thank Sam Payne for his elegant construction in Section 8.4(2).

in a transparent way. It is also natural to restrict to primitive actions. Here S_n acts *primitively* on the finite set Ω if we cannot partition Ω into disjoint blocks $\Delta_1, \Delta_2, \dots, \Delta_h$ where S_n permutes the blocks (if $\Delta_i^w \cap \Delta_j \neq \emptyset$ then $\Delta_i^w = \Delta_j$). The familiar wreath products which permute within blocks and between blocks are an example of an imprimitive action.

The primitive actions of S_n have been classified in the O’Nan-Scott theorem. We describe this carefully in Section 2. For the study of fixed points most of the cases can be handled by a marvelous theorem of Luczak-Pyber [LuPy]. This shows that, except for the action of S_n on k -sets of an n set, almost all permutations have no fixed points (we say w is a derangement). This result is explained in Section 3. For S_n acting on k -sets, one can assume that $k < n/2$, and there is a nontrivial limit if and only if k stays fixed as n tends to infinity. In these cases, the limit is shown to be an explicit polynomial in independent Poisson random variables. This is the main content of Section 4. Section 5 works out precise asymptotics for the distribution of fixed points in the action of S_n on matchings. Section 6 considers more general imprimitive subgroups. Section 7 proves that the proportion of elements of S_n which belong to a primitive subgroup not containing A_n is at most $O(n^{-2/3+\alpha})$ for any $\alpha > 0$; this improves the bound of Luczak and Pyber [LuPy]. Finally, Section 8 surveys related results (including analogs of our main results for finite classical groups) and applications of the distribution of fixed points and derangements.

If a finite group G acts on Ω with $F(w)$ the number of fixed points of w , the “lemma that is not Burnside’s” implies that

$$\begin{aligned} E(F(w)) &= \# \text{ orbits of } G \text{ on } \Omega \\ E(F^2(w)) &= \# \text{ orbits of } G \text{ on } \Omega \times \Omega = \text{rank} := r. \end{aligned}$$

If G is transitive on Ω with isotropy group H , then the rank is also the number of orbits of H on Ω and so equal to the number of $H - H$ double cosets in G .

Thus for transitive actions

$$(1.2) \quad E(F(w)) = 1, \quad \text{Var}(F(w)) = \text{rank} - 1$$

In most of our examples $P(F(w) = 0) \rightarrow 1$ but because of (2), this cannot be seen by the method moments. The standard second moment method (Durrett [Du], page 16) says that a non-negative integer random variable satisfies $P(X > E(X)/2) \geq \frac{1}{4}(EX)^2/E(X^2)$. Specializing to our case, $P(F(w) > 0) \geq 1/(4r)$; thus $P(F(w) = 0) \leq 1 - 1/(4r)$. This shows that the convergence to 1 cannot be too rapid.

There is also a quite easy lower bound for $P(F(w) = 0)$ [GW]. Even the simplest instance of this lower bound was only observed in 1992 in [CaCo].

We record this.

Theorem 1.3. *Let G be a finite transitive permutation group of degree n and rank r . Then*

$$\frac{r-1}{n} \leq P(F(w) = 0) \leq 1 - \frac{1}{4r}.$$

Frobenius groups of order $n(n-1)$ with n a prime power are only the possibilities when the lower bound is achieved. The inequality above shows that $P(F(w) = 0)$ tends to 1 implies that the rank tends to infinity. Indeed, for primitive actions of symmetric and alternating groups, this is also a sufficient condition – see Theorem 3.3.

2. O’NAN-SCOTT THEOREM

Let G act transitively on the finite set Ω . By standard theory we may represent $\Omega = G/G_\alpha$, with any fixed $\alpha \in \Omega$. Here $G_\alpha = \{w : \alpha^w = \alpha\}$ with the action being left multiplication on the cosets. Further (Passman [P, 3.4]) the action of G on Ω is primitive if and only if the isotropy group G_α is maximal. Thus, classifying primitive actions of G is the same problem as classifying maximal subgroups H of G .

The O’Nan-Scott theorem classifies maximal subgroups of A_n and S_n up to determining the almost simple primitive groups of degree n .

Theorem 2.1. [O’Nan-Scott] *Let H be a maximal subgroup of $G = A_n$ or S_n . Then, one of the following three cases holds:*

- I:** H acts primitively as a subgroup of S_n (primitive case),
- II:** $H = (S_a \wr S_b) \cap G$ (wreath product), $n = a \cdot b$, $|\Omega| = a^b \cdot b!$ (imprimitive case), or
- III:** $H = (S_k \times S_{n-k}) \cap G$, $|\Omega| = \binom{n}{k}$ with $1 \leq k < n/2$ (intransitive case).

Further, in case I, one of the following holds:

- Ia:** H is almost simple,
- Ib:** H is diagonal,
- Ic:** H preserves product structure, or
- Id:** H is affine.

Remarks and examples:

- (1) Note that in cases I, II, III, the modifiers ‘primitive’, ‘imprimitive’, ‘intransitive’ apply to H . Since H is maximal in G , $\Omega \cong G/H$ is a primitive G -set. We present an example and suitable additional definitions for each case.
- (2) In case III, Ω is the k -sets of $\{1, 2, \dots, n\}$ with the obvious action of S_n . This case is discussed extensively in Section 4 below.
- (3) In case II, take n even with $a = 2, b = n/2$. We may identify Ω with the set of perfect matchings on n points – partitions of n into $n/2$ two-element subsets where order within a subset or among subsets does not matter. For example if $n = 6$, $\{1, 2\}\{3, 4\}\{5, 6\}$ is a perfect

matching. For this case, $|\Omega| = \frac{n!}{2^{n/2}(n/2)!} = (2n-1)(2n-3)\cdots(1)$.

Careful asymptotics for this case are developed in Section 5. More general imprimitive subgroups are considered in Section 6.

- (4) While every maximal subgroup of A_n or S_n falls into one of the categories of the O’Nan-Scott theorem, not every group is maximal. A complete list of the exceptional examples is in Liebeck, Praeger and Saxl [LPS1].
- (5) In case Ia, H is *almost simple* if for some non-abelian simple group G , $G \leq H \leq \text{Aut}(G)$. For example, fix $1 < k < m$. Let $n = \binom{m}{k}$. Let S_n be all $n!$ permutations of the k sets of $\{1, 2, \dots, m\}$. Take $S_m \leq S_n$ acting on the k -sets in the usual way. For $m \geq 5$, S_m is almost simple and primitive. Here $\Omega = S_n/S_m$ does not have a simple combinatorial description, but this example is crucial and the $k = 2$ case will be analyzed in Section 7.

Let $\tau \in S_m$ be a transposition. Then τ moves precisely $2\binom{m-2}{k-1}$ elements of Ω . Thus, S_m embeds in A_n if and only if $\binom{m-2}{k-1}$ is even. Indeed for most primitive embeddings of S_m into S_n , the image is contained in A_n [NB].

It is not difficult to see that the image of S_m is maximal in either A_n or S_n . This follows from the general result in [LPS1]. It also follows from the classification of primitive groups containing a non-trivial element fixing at least $n/2$ points [GM].

Similar examples can be constructed by looking at the action of $P\Gamma L_d(q)$ on k -spaces (recall the $P\Gamma L_d(q)$ is the projective group of all semilinear transformations of a d dimensional vector space over \mathbb{F}_q). All of these are covered by case Ia.

- (6) In case Ib, H is *diagonal* if $H = G^k \dot{=} (\text{Out}(G) \times S_k)$ for G a non-abelian simple group, $k \geq 2$ (the dot denotes semidirect product). Let $\Omega = G^k/D$ with $D = \{(g, g, \dots, g)\}_{g \in G}$ the diagonal subgroup. Clearly G^k acts on Ω . Let $\text{Out}(G)$ (the outer automorphisms) act coordinate-wise and let S_k act by permuting coordinates. These transformations determine a permutation group H on the set Ω . The group H has normal subgroup G^k with quotient isomorphic to $\text{Out}(G) \times S_k$. The extension usually splits (but it doesn’t always split).

Here is an specific example. Take $G = A_m$ for $m \geq 8$ and $k = 2$. Then $\text{Out}(A_m) = C_2$ and so $H = \langle A_m \times A_m, \tau, (s, s) \rangle$ where s is a transposition (or any element in S_m outside of A_m) and τ is the involution changing coordinates. More precisely, each coset of D has a unique representative of the form $(1, x)$. We have $(g_1, g_2)(1, x)D = (g_1, g_2x)D = (1, g_2xg_1^{-1})D$. The action of $\tau \in C_2$ takes $(1, x) \rightarrow (1, x^{-1})$ and the action of $(s, s) \in \text{Out}(A_m)$ takes $(1, x)$ to $(1, sxs^{-1})$.

The maximality of H is somewhat subtle. We first show that if $m \geq 8$, then H is contained in $\text{Alt}(\Omega)$. Clearly $A_m \times A_m$ is contained

in $\text{Alt}(\Omega)$. Observe that (s, s) is contained in $\text{Alt}(\Omega)$. Indeed, taking s to be a transposition, the number of fixed points of (s, s) is the size of its centralizer in A_m which is $|S_{m-2}|$, and so $\frac{m!}{2} - (m-2)!$ points are moved and this is divisible by 4 since $m \geq 8$. To see that τ is contained in $\text{Alt}(X)$ for $m \geq 8$, note that the number of fixed points of τ is the number of involutions (including the identity) in A_m , so it is sufficient to show that $\frac{m!}{2}$ minus this number is a multiple of 4. This follows from the next proposition, which is of independent combinatorial interest.

Proposition 2.2. *Suppose that $m \geq 8$. Then the number of involutions in A_m and the number of involutions in S_m are multiples of 4.*

Proof. Let $a(m)$ be the number of involutions in A_m (including the identity). Let $b(m)$ be the number of involutions in $S_m - A_m$. It suffices to show that $a(m) = b(m) = 0 \pmod{4}$. For $n = 8, 9$ we compute directly. For $n > 9$, we observe that

$$a(n) = a(n-1) + (n-1)b(n-2)$$

and

$$b(n) = b(n-1) + (n-1)a(n-2)$$

(because an involution either fixes 1 giving the first term or swaps 1 with $j > 1$, giving rise to the second term). The result follows by induction. \square

Having verified that H is contained in A_m for $m \geq 8$, maximality now follows from Liebeck-Praeger-Saxl [LPS1].

Now consider $G = PSL(2, p)$ with $p > 11$ acting on the $n := p+1$ subspaces of dimension 1. Assume that $8 \mid (p-1)$. We claim that the number of involutions in G is always 2 modulo 4 (whence the group H described above does not embed in A_n). The centralizer of an element of order 2 has order $p-1$ and since all involutions are conjugate, there are $1 + p(p+1)/2$ involutions in G , proving the claim.

- (7) In case Ic, H preserves a product structure. Let $\Gamma = \{1, \dots, m\}, \Delta = \{1, \dots, t\}$, and let Ω be the t -fold Cartesian product of Γ . If C is a permutation group on Γ and D is a permutation group on Δ , we may define a group $H = C \wr D$ by having C act on the coordinates, and having D permute the coordinates. Primitivity of H is equivalent to C acting primitively on Γ with some non identity element having a fixed point and D acting transitively on Δ (see, e.g. Cameron [Ca1], Th. 4.5).

There are many examples of case Ic but $|\Omega| = m^t$ is rather restricted and H has a simple form. One specific example is as follows: $G = S_{m^t}$, $H = S_m \wr S_t$ and Ω is the t -fold Cartesian product

$\{1, \dots, m\}^t$. The case $t = 2$ will be analyzed in detail in Section 7. It is easy to determine when H embeds in A_{m^t} . We just note that if $t = 2$, then this is case if and only if $4|m$.

- (8) In case $\text{Id } H$ is affine. Thus $\Omega = V$, a vector space of dimension k over a field of q elements (so $n = |\Omega| = q^k$) and H is the semidirect product $V \cdot GL(V)$. Since we are interested only in maximal subgroups, q must be prime.

Note that if q is odd, then H contains an $n - 1$ cycle and so is not contained in A_n . If $q = 2$, then for $k > 2$, H is perfect and so is contained in A_n . The maximality of H in A_n or S_n follows by Mortimer [Mor] for $k > 1$ and [GKi] if $k = 1$.

- (9) The proof of the O’Nan-Scott theorem is not extremely difficult. O’Nan and Scott each presented proofs at the Santa Cruz Conference in 1979. There is a more delicate version which describes all primitive permutation groups. This was proved in Aschbacher-Scott [AS] giving quite detailed information. See also the lively lecture notes of Cameron ([Ca1], Chapter 4). The notion of generalized Fitting subgroup is quite useful in both the proof and statement of the theorem. See Kurtzweil and Stellmacher [KS].
- (10) It turns out that many of the details above are not needed for our main results. Only case III ($H = S_k \times S_{n-k}$) allows non-trivial limit theorems. This is the subject of the next section. The other cases are of interest when we try to get explicit bounds (Sections 5, 6, 7).

3. TWO THEOREMS OF LUCZAK-PYBER

The following two results are due to Luczak and Pyber.

Theorem 3.1. ([LuPy]) *Let S_n act on $\{1, 2, \dots, n\}$ as usual and let $i(n, k)$ be the number of $w \in S_n$ that leave some k -element set invariant. Then, $\frac{i(n, k)}{n!} \leq ak^{-.01}$ for an absolute constant a .*

Theorem 3.2. ([LuPy]) *Let t_n denote the number of elements of the symmetric group S_n which belong to transitive subgroups different from S_n or A_n . Then*

$$\lim_{n \rightarrow \infty} t_n/n! = 0.$$

Theorem 3.1 is at the heart of the proof of Theorem 3.2. We use them both to show that a primitive action of S_n is a derangement with probability approaching one, unless S_n acts on k -sets with fixed k . Note that we assume that $k \leq n/2$ since the action on k -sets is isomorphic to the action on $n - k$ sets.

Theorem 3.3. *Let G_i be a finite symmetric or alternating group of degree n_i acting primitively on a finite set Ω_i of cardinality at least 3. Assume that $n_i \rightarrow \infty$. Let d_i be the proportion of $w \in G_i$ with no fixed points. Then the following are equivalent:*

- (1) $\lim_{i \rightarrow \infty} d_i = 1$,
- (2) *there is no fixed k with $\Omega_i = \{k\text{-sets of } \{1, 2, \dots, n_i\}\}$ for infinitely many i , and*
- (3) *the rank of G_i acting on Ω_i tends to ∞ .*

Proof. Let H_i be an isotropy group for G_i acting on Ω_i . If H_i falls into category I or II of the O’Nan-Scott theorem, H_i is transitive. Writing out Theorem 3.2 above more fully, Luczak-Pyber prove that

$$\frac{|\bigcup_H H|}{n!} \rightarrow 0$$

where the union is over *all* transitive subgroups of S_n not equal to S_n or A_n . Thus a randomly chosen $w \in S_n$ is not in $xH_n x^{-1}$ for any x if H_n falls into category I or II.

Having ruled out categories I and II, we turn to category III (k -sets of an n set). Here, Theorem 3.1 shows the chance of a derangement tends to one as $a/k^{.01}$, for an absolute constant a .

The previous paragraphs show that (2) implies (1). If the rank does not go to ∞ , then d_i cannot approach 1 by Theorem 1.3. Thus (1) implies (3), and also (2) since the rank of the action on k -sets is $k+1$. Clearly (3) implies (2), completing the proof. \square

4. k -SETS OF AN n -SET

In this section the limiting distribution of the number of fixed points of a random permutation acting on k -sets of an n -set is determined.

Theorem 4.1. *Fix k and let S_n act on $\Omega_{n,k}$ – the k sets of $\{1, 2, \dots, n\}$. Let $A_i(w)$ be the number of i -cycles of $w \in S_n$ in its usual action on $\{1, 2, \dots, n\}$. Let $F_k(w)$ be the number of fixed points of w acting on $\Omega_{n,k}$. Then*

$$(4.2) \quad F_k(w) = \sum_{|\lambda|=k} \prod_{i=1}^k \binom{A_i(w)}{\alpha_i(\lambda)}.$$

Here the sum is over partitions λ of k and $\alpha_i(\lambda)$ is the number of parts of λ equal to i .

(2) *For all $n \geq 2$, $E(F_k) = 1$, $\text{Var}(F_k) = k$.*

(3) *As n tends to infinity, $A_i(w)$ converge to independent Poisson $(1/i)$ random variables.*

Proof. If $w \in S_n$ is to fix a k set, the cycles of w must be grouped to partition k . The expression for F_k just counts the distinct ways to do this. See the examples below. This proves (1). The rank of S_n acting on k sets is $k+1$, proving (2).

The joint limiting distribution of the A_i is a classical result due to Goncharov [Go]. In fact, letting X_1, X_2, \dots, X_k be independent Poisson with

parameters $1, \frac{1}{2}, \dots, \frac{1}{k}$, one has from [DS] that for all $n \geq \sum_{i=1}^k ib_i$,

$$E \left(\prod_{i=1}^k A_i(w)^{b_i} \right) = \prod_{i=1}^k E(X_i^{b_i}).$$

For total variation bounds see [AT]. This proves (3). \square

Examples. Throughout, let X_1, X_2, \dots, X_k be independent Poisson random variables with parameters $1, 1/2, 1/3, \dots, 1/k$ respectively.

$k = 1$: This is the usual action of S_n on $\{1, 2, \dots, n\}$ and Theorem 4.1 yields (1) of the introduction: In particular, for derangements

$$P(F_1(w) = 0) \rightarrow 1/e \doteq .36788.$$

$k = 2$: Here $F_2(w) = \binom{A_1(w)}{2} + A_2(w)$ and Theorem 4.1 says that

$$P(F_2(w) = j) \rightarrow P \left(\binom{X_1}{2} + X_2 = j \right) \text{ with } X_1 \text{ Poisson}(1), X_2 \text{ Poisson}(\tfrac{1}{2}).$$

In particular

$$P(F_2(w) = 0) \rightarrow \frac{2}{e^{3/2}} \doteq .44626.$$

$k = 3$: Here $F_3(w) = \binom{A_1(w)}{3} + A_1(w)A_2(w) + A_3(w)$ and

$$P(F_3(w) = j) \rightarrow P \left(\binom{X_1}{3} + X_1X_2 + X_3 = j \right).$$

In particular

$$P(F_3(w) = 0) \rightarrow \frac{1}{e^{4/3}} (1 + \frac{3}{2}e^{-1/2}) \doteq .50342.$$

We make the following conjecture, which has also been independently stated as a problem by Cameron [Ca2].

Conjecture: $\lim_{n \rightarrow \infty} P(F_k(w) = 0)$ is increasing in k .

Using Theorem 4.1, one can prove the following result which improves, in this context, the upper bound given in Theorem 1.3.

Proposition 4.3.

$$\lim_{n \rightarrow \infty} P(F_k(w) = 0) \leq 1 - \frac{\log(k)}{k} + O\left(\frac{1}{k}\right).$$

Proof. Clearly

$$\begin{aligned} P(F_k(w) > 0) &\geq P \left(\bigcup_{j=1}^{\lfloor \frac{k-1}{2} \rfloor} (A_{k-j} > 0 \text{ and } A_j > 0) \right) \\ &= 1 - P \left(\bigcap_{j=1}^{\lfloor \frac{k-1}{2} \rfloor} (A_{k-j} > 0 \text{ and } A_j > 0) \right). \end{aligned}$$

By Theorem 4.1, this converges to

$$1 - \prod_{j=1}^{\lfloor \frac{k-1}{2} \rfloor} \left[1 - (1 - e^{-1/j})(1 - e^{-1/(k-j)}) \right].$$

Let $a_j = (1 - e^{-1/j})$. Write the general term in the product as $e^{\log(1 - a_j a_{k-j})}$. Expand the log to $-a_j a_{k-j} + O((a_j a_{k-j})^2)$. Writing $a_j = \left(\frac{1}{j} + O\left(\frac{1}{j^2}\right)\right)$ and multiplying out, we must sum

$$\sum_{j=1}^{\lfloor \frac{k-1}{2} \rfloor} \frac{1}{j} \frac{1}{k-j}, \quad \sum_{j=1}^{\lfloor \frac{k-1}{2} \rfloor} \frac{1}{j^2} \frac{1}{k-j}, \quad \sum_{j=1}^{\lfloor \frac{k-1}{2} \rfloor} \frac{1}{(k-j)^2} \frac{1}{k}, \quad \sum_{j=1}^{\lfloor \frac{k-1}{2} \rfloor} \frac{1}{(k-j)^2} \frac{1}{j^2}.$$

Writing $\frac{1}{j} \frac{1}{k-j} = \frac{1}{k} \left(\frac{1}{j} + \frac{1}{k-j} \right)$, the first sum is $\frac{\log(k)}{k} + O\left(\frac{1}{k}\right)$. The second sum is $O\left(\frac{1}{k}\right)$, the third sum is $O\left(\frac{\log(k)}{k^2}\right)$ and the fourth is $O\left(\frac{1}{k^2}\right)$. Thus $-a_j a_{k-j}$ summed over $1 \leq j \leq (k-1)/2$ is $-\frac{\log(k)}{k} + O\left(\frac{1}{k}\right)$. The sum of $(a_j a_{k-j})^2$ is of lower order by similar arguments. In all, the lower bound on $\lim_{n \rightarrow \infty} P(F_k(w) > 0)$ is

$$1 - e^{-\frac{\log(k)}{k} + O\left(\frac{1}{k}\right)} = \frac{\log(k)}{k} + O\left(\frac{1}{k}\right).$$

□

To close this section, we give a combinatorial interpretation for the moments of the numbers $F_k(w)$ of Theorem 4.1 above. This involves the “top k to random” shuffle, which removes k cards from the top of the deck, and randomly interleaves them with the other n-k cards (choosing one of the $\binom{n}{k}$ possible interleavings uniformly at random).

Proposition 4.4. (1) *The eigenvalues of the top k to random shuffle*

are the numbers $\left\{ \frac{F_k(w)}{\binom{n}{k}} \right\}$, where w ranges over S_n .

(2) *For all values of n, k, r , the r th moment of the distribution of fixed k -sets is equal to $\binom{n}{k}^r$ multiplied by the chance that the top k to random shuffle is at the identity after r steps.*

Proof. Note that the top k to random shuffle is the inverse of the move k to front shuffle, which picks k cards at random and moves them to the front of the deck, preserving their relative order. Hence their transition matrices are transposes, so have the same eigenvalues. The move k to front shuffle is a special case of the theory of random walk on chambers of hyperplane arrangements developed in [BHR]. The arrangement is the braid arrangement and one assigns weight $\frac{1}{\binom{n}{k}}$ to each of the block ordered partitions where the first block has size k and the second block has size $n - k$. The result now follows from Corollary 2.2 of [BHR], which determined the eigenvalues of such hyperplane walks.

For the second assertion, let M be the transition matrix for the top k to random shuffle. Clearly $\text{Tr}(M^r)$ (the trace of M^r) is equal to $n!$ multiplied by the chance that the top k to random shuffle is at the identity after r steps. The first part gives that

$$\text{Tr}(M^r) = \sum_{w \in S_n} \left(\frac{F_k(w)}{\binom{n}{k}} \right)^r,$$

which implies the result. \square

As an example of part 2 of Proposition 4.4, the chance of being at the identity after 1 step is $\frac{1}{\binom{n}{k}}$ and the chance of being at the identity after 2 steps is $\frac{k+1}{\binom{n}{k}^2}$, giving another proof that $E(F_k(w)) = 1$ and $E(F_k^2(w)) = k+1$.

Remarks

- (1) As in the proof of Proposition 4.1, the moments of $F_k(w)$ can be expressed exactly in terms of the moments of Poisson random variables, provided that n is sufficiently large.
- (2) There is a random walk on the irreducible representations of S_n which has the same eigenvalues as the top k to random walk, but with different multiplicities. Unlike the top k to random walk, this walk is reversible with respect to its stationary distribution, so that spectral techniques (and hence information about the distribution of fixed points) can be used to analyze its convergence rate. For details, applications, and a generalization to other actions, see [F1], [F2].

5. FIXED POINTS ON MATCHINGS

Let M_{2n} be the set of perfect matchings on $2n$ points. Thus, if $2n = 4$, $M_{2n} = \{(1, 2)(3, 4), (1, 3)(2, 4), (1, 4)(2, 3)\}$. It is well known that

$$|M_{2n}| = (2n - 1)!! = (2n - 1)(2n - 3) \cdots (3)(1).$$

The literature on perfect matchings is enormous. See Lovász and Plummer [LoPl] for a book length treatment. Connections with phylogenetic trees and further references are in [DH1, DH2]. As explained above, the symmetric group S_{2n} acts primitively on M_{2n} . Results of Luczak-Pyber [LuPy] imply that, in this action, almost every permutation is a derangement. In this section we give sharp asymptotic rates for this last result. We show that the proportion of derangements in S_{2n} is

$$(5.1) \quad 1 - \frac{A(1)}{\sqrt{\pi n}} + o\left(\frac{1}{\sqrt{n}}\right), \quad A(1) = \prod_{i=1}^{\infty} \cosh(1/(2i - 1)).$$

Similar asymptotics are given for the proportion of permutations with $j > 0$ fixed points. This is zero if j is even. For odd j , it is

$$(5.2) \quad \frac{C(j)B(1)}{\sqrt{\pi n}} + o\left(\frac{1}{\sqrt{n}}\right), \quad B(1) = \prod_{i=1}^{\infty} \left(1 + \frac{1}{2i}\right) e^{-1/2i}$$

and $C(j)$ explicit rational numbers. In particular

$$(5.3) \quad C(1) = \frac{3}{2}, \quad C(3) = \frac{1}{4}, \quad C(5) = \frac{27}{400}, \quad C(7) = \frac{127}{2352}.$$

The argument proceeds by finding explicit closed forms for generating functions followed by standard asymptotics. It is well known that the rank of this action is $p(n)$, the number of partitions of n . Thus (5.1) is a big improvement over the upper bound given in Theorem 1.3.

For $w \in S_{2n}$, let $a_i(w)$ be the number of i -cycles in the cycle decomposition. Let $F(w)$ be the number of fixed points of w acting on M_{2n} . The following proposition determines $F(w)$ in terms of $a_i(w)$, $1 \leq i \leq 2n$.

Proposition 5.4. *The number of fixed points, $F(w)$ of $w \in S_{2n}$ on M_{2n} is*

$$F(w) = \prod_{i=1}^{2n} F_i(a_i(w))$$

with

$$F_{2i-1}(a) = \begin{cases} 1 & \text{if } a = 0 \\ 0 & \text{if } a \text{ is odd} \\ (a-1)!!(2i-1)^{a/2} & \text{if } a > 0 \text{ is even} \end{cases}$$

$$F_{2i}(a) = 1 + \sum_{k=1}^{\lfloor a/2 \rfloor} (2k-1)!! \binom{a}{2k} (2i)^k$$

In particular,

$$(5.5) \quad F(w) \neq 0 \text{ if and only if } a_{2i-1}(w) \text{ is even for all } i$$

$$(5.6) \quad F(w) \text{ does not take on non-zero even values.}$$

Proof. Consider first the cycles of w of length $2i-1$. If a_{2i-1} is even, the cycles may be matched in pairs, then each pair of length $2i-1$ can be broken into two element matched two element subsets by first pairing the lowest element among the numbers with any of the $2i-1$ numbers in the second cycle. The rest is determined by cyclic action. For example, if the two three-cycles (123)(456) appear, the matched pairs (14)(25)(36) are fixed, so are (15)(26)(34) or (16)(24)(35). Thus $F_3(2) = 3$. If a_{2i-1} is odd, some element cannot be matched and $F_{2i-1}(a_{2i-1}) = 0$.

Consider next the cycles of w of length $2i$. Now, there are two ways to create parts of a fixed perfect matching. First, some of these cycles can be paired and, for each pair, the previous construction can be used. Second, for each unpaired cycle, elements i apart can be paired. For example,

from (1234) the pairing (13)(24) may be formed. The sum in $F_{2i}(a)$ simply enumerates by partial matchings.

To see that $F(w)$ cannot take on non-zero even values, observe that $F_{2i-1}(a)$ and $F_{2i}(a)$ only take on odd values if they are non-zero. \square

Let $P_{2n}(j) = \frac{|\{w \in S_{2n} : F(w) = j\}|}{2n!}$. For $j \geq 1$, let $g_j(t) = \sum_{n=0}^{\infty} t^{2n} P_{2n}(j)$ and let $\bar{g}_0(t) = \sum_{n=0}^{\infty} t^{2n} (1 - P_{2n}(0))$.

Proposition 5.7.

$$\bar{g}_0(t) = \frac{\prod_{i=1}^{\infty} \cosh(t^{2i-1}/(2i-1))}{\sqrt{1-t^2}} \quad \text{for } 0 < t < 1.$$

Proof. From Proposition 5.4, $w \in S_{2n}$ has $F(w) \neq 0$ if and only if $a_{2i-1}(w)$ is even for all i . From Shepp-Lloyd [SL], if N is chosen in $\{0, 1, 2, \dots\}$ with

$$P(N = n) = (1-t)t^n$$

and then w is chosen uniformly in S_N , the $a_i(w)$ are independent Poisson random variables with parameter t^i/i respectively. If X is a Poisson (λ) random variable, $P(X \text{ is even}) = \left(\frac{1}{2} + \frac{e^{-2\lambda}}{2}\right)$. It follows that

$$\begin{aligned} \sum_{n=0}^{\infty} (1-t)t^{2n} (1 - P_{2n}(0)) &= \prod_{i=1}^{\infty} \left(\frac{1 + e^{-2t^{2i-1}/(2i-1)}}{2} \right) \\ &= \prod_{i=1}^{\infty} e^{-t^{2i-1}/(2i-1)} \prod_{i=1}^{\infty} \cosh\left(\frac{t^{2i-1}}{(2i-1)}\right) \\ &= \sqrt{\frac{1-t}{1+t}} \prod_{i=1}^{\infty} \cosh(t^{2i-1}/(2i-1)). \end{aligned}$$

\square

Corollary 5.8. *As n tends to infinity,*

$$1 - P_{2n}(0) \sim \frac{\prod_{i=1}^{\infty} \cosh(1/(2i-1))}{\sqrt{\pi n}}.$$

Proof. By Proposition 5.7, $1 - P_{2n}(0)$ is the coefficient of t^n in

$$\frac{\prod_{i=1}^{\infty} \cosh(t^{(2i-1)/2}/(2i-1))}{\sqrt{1-t}}.$$

It is straightforward to check that the numerator is analytic near $t = 1$, so the result follows from Darboux's theorem ([O], Theorem 11.7). \square

Proposition 5.4 implies that the event $F(w) = j$ is contained in the event $\{a_{2i-1}(w) \text{ is even for all } i \text{ and } a_{2i}(w) \in \{0, 1\} \text{ for } 2i > j\}$. This is evidently complicated for large j .

We prove

Proposition 5.9. *For positive odd j ,*

$$g_j(t) = \frac{P_j(t) \prod_{i=1}^{\infty} \left(1 + \frac{t^{2i}}{2i}\right) e^{-t^{2i}/2i}}{\sqrt{1-t^2}}$$

for $P_j(t)$ an explicit rational function in t with positive rational coefficients. In particular,

$$\begin{aligned} P_1(t) &= \left(1 + \frac{t^2}{2}\right), P_3(t) = \left(\frac{t^4}{6} + \frac{t^6}{18} + \frac{t^8}{36}\right) \\ P_5(t) &= \frac{\frac{1}{2}\left(1 + \frac{t^2}{2}\right)\left(\frac{t^2}{4}\right)^2}{1 + \frac{t^4}{4}} + \frac{1}{2}\left(1 + \frac{t^2}{2}\right)\left(\frac{t^5}{5}\right)^2 \\ P_7(t) &= \frac{\frac{1}{2}\left(1 + \frac{t^2}{2}\right)\left(\frac{t^6}{6}\right)^2}{1 + \frac{t^6}{6}} + \frac{1}{6}\left(\frac{t^2}{2}\right)^3 + \frac{1}{2}\left(1 + \frac{t^2}{2}\right)\left(\frac{t^7}{7}\right)^2. \end{aligned}$$

Proof. Consider first the case of $j = 1$. From Proposition 5.4, $F(w) = 1$ if and only if $a_{2i-1}(w) = 0$ for $i \geq 2$, $a_1(w) \in \{0, 2\}$ and $a_{2i}(w) \in \{0, 1\}$ for all $i \geq 1$. For example, if $2n = 10$ and $w = (1)(2)(345678910)$, the unique fixed matching is $(1\ 2)(3\ 7)(4\ 8)(5\ 9)(6\ 10)$. From the cycle index argument used in Proposition 5.7,

$$\begin{aligned} & \sum_{n=0}^{\infty} (1-t)t^{2n} P_{2n}(F(w) = 1) \\ &= e^{-t} \left(1 + \frac{t^2}{2}\right) \prod_{i=2}^{\infty} e^{-t^{2i-1}/(2i-1)} \prod_{i=1}^{\infty} e^{-t^{2i}/2i} \left(1 + \frac{t^{2i}}{2i}\right) \\ &= (1-t) \left(1 + \frac{t^2}{2}\right) \prod_{i=1}^{\infty} \left(1 + \frac{t^{2i}}{2i}\right) \\ &= \frac{(1-t) \left(1 + \frac{t^2}{2}\right) \prod_{i=1}^{\infty} \left(1 + \frac{t^{2i}}{2i}\right) e^{-t^{2i}/2i}}{\sqrt{1-t^2}}. \end{aligned}$$

The arguments for the other parts are similar. In particular, $F(w) = 3$ iff one of the following holds:

- $a_1(w) = 4$, all $a_{2i-1}(w) = 0$ $i \geq 2$, all $a_{2i}(w) \in \{0, 1\}$
- $a_1(w) \in \{0, 2\}$, $a_2(w) = 2$ all $a_{2i-1}(w) = 0$ and $a_{2i}(w) \in \{0, 1\}$ $i \geq 2$
- $a_1(w) \in \{0, 2\}$, $a_3(w) = 2$, $a_{2i-1}(w) = 0$ $i \geq 3$, $a_{2i} \in \{0, 1\}$

Similarly, $F(w) = 5$ iff one of the following holds:

- $a_4(w) = 2$, $a_1(w) \in \{0, 2\}$, $a_{2i-1}(w) = 0$, $a_{2i}(w) \in \{0, 1\}$ else
- $a_5(w) = 2$, $a_1(w) \in \{0, 2\}$, $a_{2i-1}(w) = 0$, $a_{2i}(w) \in \{0, 1\}$ else

Finally, $F(w) = 7$ iff one of the following holds:

- $a_1(w) \in \{0, 2\}$, $a_6(w) = 2$ or $a_2(w) = 3$, $a_{2i-1}(w) = 0$, $a_{2i}(w) \in \{0, 1\}$ else
- $a_7(w) = 2$, $a_1(w) \in \{0, 1\}$, $a_{2i-1}(w) = 0$, $a_{2i}(w) \in \{0, 1\}$ else

Further details are omitted. \square

The asymptotics in (2) follow from Proposition 5.9, by the same method used to prove (1) in Corollary 5.8.

6. MORE IMPRIMITIVE SUBGROUPS

Section 5 studied fixed points on matchings, or equivalently fixed points of S_{2n} on the left cosets of $S_2 \wr S_n$. This section uses a quite different approach to study derangements of S_{an} on the left cosets of $S_a \wr S_n$, where $a \geq 2$ is constant. It is proved that the proportion of elements of S_{an} which fix at least one left coset of $S_a \wr S_n$ (or equivalently are conjugate to an element of $S_a \wr S_n$ or equivalently fix a system of n blocks of size a) is at most the coefficient of u^n in

$$\exp \left(\sum_{k \geq 1} \frac{u^k}{a!} \binom{1}{k} \left(\frac{1}{k} + 1\right) \cdots \left(\frac{1}{k} + a - 1\right) \right),$$

and that this coefficient is asymptotic to $C_a n^{\frac{1}{a}-1}$ as $n \rightarrow \infty$, where C_a is an explicit constant depending on a (defined in Proposition 6.3 below). In the special case of matchings ($a = 2$), this becomes $\frac{e^{\frac{1}{2}}}{\sqrt{\pi n}}$, which is extremely close to the true asymptotics obtained in Section 5. Moreover, this generating function will be crucially applied when we sharpen a result of Luczak and Pyber in Section 7.

The method of proof is straightforward. Clearly the number of permutations in S_{an} conjugate to an element of $S_a \wr S_n$ is upper bounded by the sum over conjugacy classes C of $S_a \wr S_n$ of the size of the S_{an} conjugacy class of C . Unfortunately this upper bound is hard to compute, but we show it to be smaller than something which can be exactly computed as a coefficient in a generating function. This will prove the result.

From Section 4.2 of [JK], there is the following useful description of conjugacy classes of $G \wr S_n$ where G is a finite group. The classes correspond to matrices M with natural number entries $M_{i,k}$, rows indexed by the conjugacy classes of G , columns indexed by the numbers $1, 2, \dots, n$, and satisfying the condition that $\sum_{i,k} k M_{i,k} = n$. More precisely, given an element $(g_1, \dots, g_n; \pi)$ in $G \wr S_n$, for each k -cycle of π one multiplies the g 's whose subscripts are the elements of the cycle in the order specified by the cycle. Taking the conjugacy class in G of the resulting product contributes 1 to

the matrix entry whose row corresponds to this conjugacy class in G and whose column is k .

The remainder of this section specializes to $G = S_a$. Since conjugacy classes of S_a correspond to partitions λ of a , the matrix entries are denoted by $M_{\lambda,k}$. We write $|\lambda| = a$ if λ is a partition of a . Given a partition λ , let $n_i(\lambda)$ denote the number of parts of size i of λ .

Proposition 6.1. *Let the conjugacy class C of $S_a \wr S_n$ correspond to the matrix $(M_{\lambda,k})$ where λ is a partition of a . Then the proportion of elements of S_{an} conjugate to an element of C is at most*

$$\frac{1}{\prod_k \prod_{|\lambda|=a} M_{\lambda,k}! [\prod_i (ik)^{n_i(\lambda)} n_i(\lambda)!]^{M_{\lambda,k}}}.$$

Proof. Observe that the number of cycles of length j of an element of C is equal to

$$\sum_{k|j} \sum_{|\lambda|=a} M_{\lambda,k} n_{j/k}(\lambda).$$

To see this, note that $S_a \wr S_n$ can be viewed concretely as a permutation of an symbols by letting it act on an array of n rows of length a , with S_a permuting within each row and S_n permuting among the rows.

Hence by a well known formula for conjugacy class sizes in a symmetric group, the proportion of elements of S_{an} conjugate to an element of C is equal to

$$\begin{aligned} & \frac{1}{\prod_j j^{\sum_{k|j} \sum_{|\lambda|=a} M_{\lambda,k} n_{j/k}(\lambda)} [\sum_{k|j} \sum_{|\lambda|=a} M_{\lambda,k} n_{j/k}(\lambda)]!} \\ & \leq \frac{1}{\prod_j j^{\sum_{k|j} \sum_{|\lambda|=a} M_{\lambda,k} n_{j/k}(\lambda)} \prod_{k|j} \prod_{|\lambda|=a} M_{\lambda,k} n_{j/k}(\lambda)!} \\ & \leq \frac{1}{\prod_j j^{\sum_{k|j} \sum_{|\lambda|=a} M_{\lambda,k} n_{j/k}(\lambda)} \prod_{k|j} \prod_{|\lambda|=a} [M_{\lambda,k}! n_{j/k}(\lambda)!]^{M_{\lambda,k}}} \\ & = \frac{1}{\prod_k \prod_{|\lambda|=a} M_{\lambda,k}! [\prod_i (ik)^{n_i(\lambda)} n_i(\lambda)!]^{M_{\lambda,k}}}, \end{aligned}$$

as desired. The first inequality uses the fact that $(x_1 + \dots + x_n)! \geq x_1! \dots x_n!$. The second inequality uses that $(xy)! \geq x!y!^x$ for $x, y \geq 1$ integers, which is true since

$$(xy)! = \prod_{i=1}^x \prod_{j=0}^{y-1} (i+jx) \geq \prod_{i=1}^x \prod_{j=0}^{y-1} i(1+j) = (x!)^y (y!)^x \geq x!(y!)^x.$$

The final equality used the change of variables $i = j/k$. \square

To proceed further, the next lemma is useful.

Lemma 6.2.

$$\sum_{|\lambda|=a} \frac{1}{\prod_i (ik)^{n_i(\lambda)} n_i(\lambda)!} = \frac{(\frac{1}{k})(\frac{1}{k} + 1) \cdots (\frac{1}{k} + a - 1)}{a!}.$$

Proof. Let $c(\pi)$ denote the number of cycles of a permutation π . Since the number of permutations in S_a with n_i cycles of length i is $\frac{a!}{\prod_i i^{n_i} n_i!}$, the left hand side is equal to

$$\frac{1}{a!} \sum_{\pi \in S_a} k^{-c(\pi)}.$$

It is well known and easily proved by induction that

$$\sum_{\pi \in S_a} x^{c(\pi)} = x(x+1) \cdots (x+a-1).$$

□

Theorem 6.3 applies the preceding results to obtain a useful generating function.

Theorem 6.3. (1) *The proportion of elements in S_{an} conjugate to an element of $S_a \wr S_n$ is at most the coefficient of u^n in*

$$\exp \left(\sum_{k \geq 1} \frac{u^k}{a!} \left(\frac{1}{k}\right) \left(\frac{1}{k} + 1\right) \cdots \left(\frac{1}{k} + a - 1\right) \right).$$

(2) *For a fixed a and $n \rightarrow \infty$, the coefficient of u^n in this generating function is asymptotic to*

$$\frac{e^{\sum_{r=2}^a p(a,r)\zeta(r)}}{\Gamma(1/a)} n^{\frac{1}{a}-1}$$

where $p(a,r)$ is the proportion of permutations in S_a with exactly r cycles, ζ is the Riemann zeta function, and Γ is the gamma function.

Proof. Proposition 6.1 implies that the sought proportion is at most the coefficient of u^n in

$$\begin{aligned} & \prod_k \prod_{|\lambda|=a} \sum_{M_{\lambda,k} \geq 0} \frac{u^{kM_{\lambda,k}}}{M_{\lambda,k}! [(ik)^{n_i(\lambda)} n_i(\lambda)!]^{M_{\lambda,k}}} \\ &= \prod_k \prod_{|\lambda|=a} \exp \left(\frac{u^k}{(ik)^{n_i(\lambda)} n_i(\lambda)!} \right) \\ &= \prod_k \exp \left(\sum_{|\lambda|=a} \frac{u^k}{(ik)^{n_i(\lambda)} n_i(\lambda)!} \right) \\ &= \exp \left(\sum_{k \geq 1} \frac{u^k}{a!} \left(\frac{1}{k}\right) \left(\frac{1}{k} + 1\right) \cdots \left(\frac{1}{k} + a - 1\right) \right). \end{aligned}$$

The last equality used Lemma 6.2.

For the second assertion, one uses Darboux's lemma (see [O] for an exposition), which gives the asymptotics of functions of the form $(1-u)^\alpha g(u)$ where $g(u)$ is analytic near 1, $g(1) \neq 0$, and $\alpha \notin \{0, 1, 2, \dots\}$. More precisely it gives that the coefficient of u^n in $(1-u)^\alpha g(u)$ is asymptotic to $\frac{g(1)}{\Gamma(-\alpha)} n^{-\alpha-1}$. By Lemma 6.2,

$$\begin{aligned} & \exp \left(\sum_{k \geq 1} \frac{u^k}{a!} \left(\frac{1}{k}\right) \left(\frac{1}{k} + 1\right) \cdots \left(\frac{1}{k} + a - 1\right) \right) \\ &= \exp \left(\sum_{k \geq 1} \frac{u^k}{ak} + \sum_{k \geq 1} u^k \sum_{r=2}^a p(a, r) k^{-r} \right) \\ &= (1-u)^{-\frac{1}{a}} \cdot \exp \left(\sum_{r=2}^a p(a, r) \sum_{k \geq 1} \frac{u^k}{k^r} \right). \end{aligned}$$

Taking $g(u) = \exp \left(\sum_{r=2}^a p(a, r) \sum_{k \geq 1} \frac{u^k}{k^r} \right)$ proves the result. \square

Remark The upper bound in Theorem 6.3 is not perfect. In fact when $n = 2$, it does not approach 0 as $a \rightarrow \infty$, whereas the true answer must by Theorem 3.3. However by part 2 of Theorem 6.3, the bound is useful for a fixed and n growing, and it will be crucially applied in Section 7 when $a = n$ are both growing.

7. PRIMITIVE SUBGROUPS

A main goal of this section is to prove that the proportion of elements of S_n which belong to a primitive subgroup not containing A_n is at most $O(n^{-2/3+\alpha})$ for any $\alpha > 0$. This improves on the bound $O(n^{-1/2+\alpha})$ in [LuPy], which was used in proving Theorem 3.2 in Section 2. We conjecture that this can in fact be replaced by $O(n^{-1})$ (and the examples with $n = (q^d - 1)/(q - 1)$ with the subgroup containing $PGL(d, q)$ or $n = p^d$ with subgroup $AGL(d, p)$ show that in general one can do no better).

The minimal degree of a permutation group is defined as the least number of points moved by a nontrivial element. The first step is to classify the degree n primitive permutation groups with minimal degree at most $n^{2/3}$.

Theorem 7.1. *Let G be a primitive permutation group of degree n . Assume that there is a nontrivial $g \in G$ moving at most $n^{2/3}$ points. Then one of the following holds:*

- (1) $G = A_n$ or S_n ;
- (2) $G = S_m$ or A_m with $m \geq 5$ and $n = \binom{m}{2}$ (acting on subsets of size 2); or
- (3) $A_m \times A_m < G \leq S_m \wr S_2$ with $n = m^2$ (preserving a product structure).

Proof. First note that the minimal degree in (2) is $2(m-2)$ and in (3) is $2n^{1/2}$. In particular, aside from (1), we always have the minimal degree is at least $n^{1/2}$.

It follows by the main result of [GM] that if there is a $g \in G$ moving fewer than $n/2$ points, then one the following holds:

(a) G is almost simple with socle (the subgroup generated by the minimal normal subgroups) A_m and $n = \binom{m}{k}$ with the action on subsets of size $k < m/2$;

(b) $n = m^t$, with $t > 1$, $m \geq 5$, G has a unique minimal normal subgroup $N = L_1 \times \dots \times L_t$ with $t > 1$ and G preserves a product structure – i.e. if $\Omega = \{1, \dots, n\}$, then as G -sets, $\Omega \cong X^t$ where $m = |X|$, $G \leq S_m \wr S_t$ acts on X^t by acting on each coordinate and permuting the coordinates.

Note that $n/2 \geq n^{2/3}$ as long as $n \geq 8$. If $n < 8$, then G contains an element moving at most 3 points, i.e. either a transposition or a 3 cycle, and so contains A_n (Theorem 3.3A in [DxM]).

Consider (a) above. If $k = 1$, then (1) holds. If $3 \leq k < m/2$, then it is an easy exercise to see that the element of S_m moving the fewest k sets is a transposition. The number of k -sets moved is $2\binom{m-2}{k-1}$. We claim that this is greater than $n^{2/3}$. Indeed, the sought inequality is equivalent to checking that $\frac{2k(m-k)}{m(m-1)} > \binom{m}{k}^{-1/3}$. The worst case is clearly $k = 3$, which is checked by taking cubes. This settles the case $3 \leq k < m/2$, and if $k = 2$, we are in case (2).

Now consider (b) above. Suppose that $t \geq 3$. Then if $g \in S_m \times \dots \times S_m$ is nontrivial, it moves at least $2m^{t-1} > n^{2/3}$ many points. If $g \in S_m \wr S_t$ and is not in $S_m \times \dots \times S_m$, then up to conjugacy we may write $g = (g_1, \dots, g_t; \sigma)$ where say σ has an orbit $\{1, \dots, s\}$ with $s > 1$. Viewing our set as $A \times B$ with A being the first s coordinates, we see that g fixes at most m points on A (since there is at most one g fixed point with a given coordinate) and so on the whole space, g fixes at most $m^{t-s+1} \leq m^{t-1}$ points and so moves at least $m^t - m^{t-1}$ points. Since $t \geq 3$, this is greater than $n^{2/3}$. Summarizing, we have shown that in case (b), $t \geq 3$ leads to a contradiction.

So finally consider (b) with $t = 2$. If g is not in $S_m \times S_m$, then arguing as in the $t = 3$ case shows that g moves at least $m^2 - m$ points. This is greater than $m^{4/3} = n^{2/3}$ since $m \geq 5$, a contradiction. So write $g = (g_1, g_2) \in S_m \times S_m$ with say $g_1 \neq 1$. If g_1 moves at least d points, then g moves at least dm points. This is greater than $n^{2/3}$ unless $d \leq m^{1/3}$. By the remark at the beginning of the proof, this implies that $L = A_m$, whence (1) holds (note there is a slight subtlety here – a priori L need not be acting primitively on the first coordinate, but in fact, it does follow from the Aschbacher-O’Nan-Scott theorem [AS] that it is contained in a primitive subgroup which normalizes L). \square

Next, we focus on Case 2 of Theorem 7.1.

Lemma 7.2. *Let S_m be viewed as a subgroup of $S_{\binom{m}{2}}$ using its action on 2-sets of $\{1, \dots, m\}$. For $w \in S_m$, let $A_i(w)$ denote the number of cycles of w of length i in its usual action on $\{1, \dots, m\}$. The total number of orbits of w on 2-sets $\{j, k\}$ of symbols which are in a common cycle of w is*

$$\frac{m}{2} - \sum_{i \text{ odd}} \frac{A_i(w)}{2}.$$

Proof. First suppose that w is a single cycle of length $i \geq 2$. If i is odd, then all orbits of w on pairs of symbols in the i -cycle have length i , so the total number of orbits is $\frac{\binom{i}{2}}{i} = \frac{i-1}{2}$. If i is even, there is 1 orbit of size $\frac{i}{2}$ and all other orbits have size i , giving a total of $\frac{i}{2}$ orbits. Hence for general w , the total number of orbits on pairs of symbols in a common cycle of w is

$$\begin{aligned} \sum_{\substack{i \text{ odd} \\ i \geq 3}} A_i(w) \frac{i-1}{2} + \sum_{i \text{ even}} A_i(w) \frac{i}{2} &= \sum_{\substack{i \text{ odd} \\ i \geq 1}} A_i(w) \frac{i-1}{2} + \sum_{i \text{ even}} A_i(w) \frac{i}{2} \\ &= \frac{m}{2} - \sum_{i \text{ odd}} \frac{A_i(w)}{2}. \end{aligned}$$

□

Theorem 7.3. *Let S_m be viewed as a subgroup of S_n with $n = \binom{m}{2}$ using its action on 2-sets of $\{1, \dots, m\}$. Then the proportion of elements of S_n contained in a conjugate of S_m is at most $O\left(\frac{\log(n)}{n}\right)$.*

Proof. We claim that any element w of S_m has at least $\frac{m}{12}$ cycles when viewed as an element of S_n . Indeed, if $A_1(w) > \frac{m}{2}$, then w fixes at least $\frac{m(m-1)}{8} \geq \frac{m}{12}$ two-sets. So we suppose that $A_1(w) \leq \frac{m}{2}$. Clearly $\sum_{i \geq 3, \text{ odd}} A_i(w) \leq \frac{m}{3}$. Thus Lemma 7.2 implies that w has at least $\frac{m}{2} - \frac{m}{4} - \frac{m}{6} = \frac{m}{12}$ cycles as an element of S_n . The number of cycles of a random element of S_n has mean and variance asymptotic to $\log(n) \sim 2 \log(m)$ (and is in fact asymptotically normal) [Go]. Thus by Chebyshev's inequality, the proportion of elements in S_m with at least $\frac{m}{12}$ cycles is $O\left(\frac{\log(m)}{m^2}\right) = O\left(\frac{\log(n)}{n}\right)$, as desired. □

To analyze Case 3 of Theorem 7.1, the following bound, based on the generating function from Section 6, will be needed.

Proposition 7.4. *The proportion of elements in S_{m^2} which fix a system of m blocks of size m is $O(n^{-3/4} + \alpha)$ for any $\alpha > 0$.*

Proof. By Theorem 6.3, the proportion in question is at most the coefficient of u^m in

$$\exp\left(\sum_{k \geq 1} \frac{u^k}{mk} \left(1 + \frac{1}{k}\right) \left(1 + \frac{1}{2k}\right) \cdots \left(1 + \frac{1}{(m-1)k}\right)\right).$$

If $f(u)$ and $g(u)$ are power series in u , we write $f(u) \ll g(u)$ if the coefficient of u^n in $f(u)$ is less than or equal to the corresponding coefficient in $g(u)$, for all n . Since $\log(1+x) \leq x$ for $0 < x < 1$, one has that

$$\log \left(\prod_{i=1}^{m-1} \left(1 + \frac{1}{ik}\right) \right) \leq \sum_{i=1}^{m-1} \frac{1}{ki} \leq \frac{1}{k} (1 + \log(m-1)).$$

Thus

$$\begin{aligned} & \exp \left(\sum_{k \geq 1} \frac{u^k}{mk} \left(1 + \frac{1}{k}\right) \left(1 + \frac{1}{2k}\right) \cdots \left(1 + \frac{1}{(m-1)k}\right) \right) \\ & \ll \exp \left(\sum_{k \geq 1} \frac{u^k}{mk} e^{1/k} (m-1)^{1/k} \right) \\ & \ll e^{ue} \exp \left(\sum_{k \geq 2} \frac{u^k}{k} \sqrt{\frac{e}{m}} \right) \\ & \ll e^{ue} \exp \left(\sum_{k \geq 1} \frac{u^k}{k} \sqrt{\frac{e}{m}} \right) \\ & = e^{ue} (1-u)^{-\sqrt{\frac{e}{m}}}. \end{aligned}$$

The coefficient of u^i in $(1-u)^{-\sqrt{\frac{e}{m}}}$ is

$$\frac{1}{i!} \sqrt{\frac{e}{m}} \prod_{j=1}^{i-1} \left(\sqrt{\frac{e}{m}} + j - 1 \right) = \frac{1}{i} \sqrt{\frac{e}{m}} \prod_{j=1}^{i-1} \left(1 + \frac{1}{j} \sqrt{\frac{e}{m}} \right).$$

Since

$$\log \left(\prod_{j=1}^{i-1} \left(1 + \frac{1}{j} \sqrt{\frac{e}{m}} \right) \right) \leq \sum_{j=1}^{i-1} \frac{1}{j} \sqrt{\frac{e}{m}} \leq \sqrt{\frac{e}{m}} (1 + \log(i-1)),$$

it follows that

$$\prod_{j=1}^{i-1} \left(1 + \frac{1}{j} \sqrt{\frac{e}{m}} \right) \leq [e(i-1)]^{\sqrt{\frac{e}{m}}}.$$

This is at most a universal constant A if $0 \leq i \leq m$. Thus the coefficient of u^m in $e^{ue} (1-u)^{-\sqrt{\frac{e}{m}}}$ is at most

$$\frac{e^m}{m!} + A \sqrt{\frac{e}{m}} \sum_{i=1}^m \frac{1}{i} \frac{e^{m-i}}{(m-i)!}.$$

By Stirling's formula (page 52 of [Fe]), $m! > m^m e^{-m+1/(12m+1)} \sqrt{2\pi m}$, which implies that the first term is very small for large m . To bound the sum, consider the terms for $i \geq m^{1-\alpha}$, where $0 < \alpha < 1$. These contribute at most $\frac{Bm^\alpha}{m^{3/2}}$ for a universal constant B . The contribution of the other

terms in negligible in comparison, by Stirling's formula. Summarizing, the contribution of the sum is $O(m^{-3/2+\alpha}) = O(n^{-3/4+\alpha/2})$, as desired. \square

The following theorem gives a bound for Case 3 of Theorem 7.1.

Theorem 7.5. *Let $S_m \wr S_2$ be viewed as a subgroup of S_n with $n = m^2$ using its action on the Cartesian product $\{1, \dots, m\}^2$. Then the proportion of elements of S_n conjugate to an element of $S_m \wr S_2$ is $O(n^{-3/4+\alpha})$ for any $\alpha > 0$.*

Proof. Consider elements of $S_m \wr S_2$ of the form $(w_1, w_2; id)$. These all fix m blocks of size m in the action on $\{1, \dots, m\}^2$; the blocks consist of points with a given first coordinate. By Proposition 7.4, the proportion of elements of S_n conjugate to some $(w_1, w_2; id)$ is $O(n^{-3/4+\alpha})$ for any $\alpha > 0$.

Next, consider an element of $S_m \wr S_2$ of the form $\sigma = (w_1, w_2; (12))$. Then $\sigma^2 = (w_1 w_2, w_2 w_1; id)$. Note that $w_1 w_2$ and $w_2 w_1$ are conjugate in S_m , and let A_i denote their common number of i -cycles. Observe that if x is in an i -cycle of $w_1 w_2$, and y is in an i -cycle of $w_2 w_1$, then $(x, y) \in \Omega$ is in an orbit of σ^2 of size i . Hence the total number of orbits of σ^2 of size i is at least $\frac{(iA_i)^2}{i} \geq iA_i$. Thus the total number of orbits of σ^2 on Ω is at least $\sum_i iA_i = m$. Hence the total number of orbits of σ is at least $\frac{m}{2}$. Arguing as in the proof of Theorem 7.3, it follows that the proportion of elements of S_n conjugate to an element of the form σ is $O\left(\frac{\log(n)}{n}\right)$, and so is $O(n^{-3/4+\alpha})$ for any $\alpha > 0$. \square

Now the main result of this section can be proved.

Theorem 7.6. *The proportion of elements of S_n which belong to a primitive subgroup not containing A_n is at most $O(n^{-2/3+\alpha})$ for any $\alpha > 0$.*

Proof. Fix $\alpha > 0$. By Bovey [Bo], the proportion of elements w of S_n such that $\langle w \rangle$ has minimum degree greater than $n^{2/3}$ is $O(n^{-2/3+\alpha})$. Thus the proportion of $w \in S_n$ which lie in a primitive permutation group having minimal degree greater than $n^{2/3}$ is $O(n^{-2/3+\alpha})$. The only primitive permutation groups of degree n with minimal degree $\leq n^{2/3}$, and not containing A_n are given by Cases 2 and 3 of Theorem 7.1. Theorems 7.3 and 7.5 imply that the proportion of w lying in the union of all such subgroups is $O(n^{-2/3+\alpha})$, so the result follows. \square

A trivial corollary to the theorem is that this holds for A_n as well.

Remark The actions of the symmetric group studied in this section embed the group as a subgroup of various larger symmetric groups. Any such embedding can be thought of as a code in the larger symmetric group. Such codes may be used for approximating sums of various functions over the larger symmetric group via a sum over the smaller symmetric group. Our results can be interpreted as giving examples of functions where the approximation is not particularly accurate. For example, the proof of Theorem 7.3

shows this to be the case when S_m is viewed as a subgroup of $S_n, n = \binom{m}{2}$ using the actions on 2-sets, and the function is the number of cycles.

8. RELATED RESULTS AND APPLICATIONS

There are numerous applications of the distribution of fixed points and derangements. Subsection 8.1 mentions some motivation from number theory. Subsection 8.2 discusses some literature on the proportion of derangements and an analog of the main result of our paper for finite classical groups. Subsection 8.3 discusses fixed point ratios, emphasizing the application to random generation. Subsection 8.4 collects some miscellany about fixed points and derangements, including algorithmic issues and appearances in algebraic combinatorics.

While this section does cover many topics, the survey is by no means comprehensive. Some splendid presentations of several other topics related to derangements are Serre [Se], Cameron's lecture notes [Ca2] and Section 6.6 of [Ca1]. For the connections with permutations with restricted positions and rook polynomials see [St, 2.3, 2.4].

8.1. Motivation from number theory. We describe two number theoretic applications of derangements which can be regarded as motivation for their study:

(1) *Zeros of polynomials* Let $h(T)$ be a polynomial with integer coefficients which is irreducible over the integers. Let $\pi(x)$ be the number of primes $\leq x$ and let $\pi_h(x)$ be the number of primes $\leq x$ for which h has no zeros mod p . It follows from Chebotarev's density theorem (see [LS] for history and a proof sketch), that $\lim_{x \rightarrow \infty} \frac{\pi_h(x)}{\pi(x)}$ is equal to the proportion of derangements in the Galois group G of $h(T)$ (viewed as permutation of the roots of $h(T)$). Several detailed examples are worked out in Serre's survey [Se].

In addition, there are applications such as the the number field sieve for factoring integers (Section 9 of [BLP]), where it is important to understand the proportion of primes for which h has no zeros mod p . This motivated Lenstra (1990) to pose the question of finding a good lower bound for the proportion of derangements of a transitive permutation group acting on a set of n letters with $n \geq 2$. Results on this question are described in Subsection 8.2.

(2) *The value problem* Let \mathbb{F}_q be a finite field of size q with characteristic p and let $f(T)$ be a polynomial of degree $n > 1$ in $\mathbb{F}_q[T]$ which is not a polynomial in T^p . The arithmetic question raised by Chowla [Ch] is to estimate the number V_f of distinct values taken by $f(T)$ as T runs over \mathbb{F}_q .

There is an asymptotic formula for V_f in terms of certain Galois groups and derangements. More precisely, let G be the Galois group of $f(T) - t = 0$ over $\mathbb{F}_q(t)$ and let N be the Galois group of $f(T) - t = 0$ over $\overline{\mathbb{F}_q}(t)$, where $\overline{\mathbb{F}_q}$ is an algebraic closure of \mathbb{F}_q . Both groups act transitively on the n roots

of $f(T) - t = 0$. The geometric monodromy group N is a normal subgroup of the arithmetic monodromy group G . The quotient group G/N is a cyclic group (possibly trivial).

Theorem 8.1. ([Co]) *Let xN be the coset which is the Frobenius generator of the cyclic group G/N . The Chebotarev density theorem for function fields yields the following asymptotic formula:*

$$V_f = \left(1 - \frac{|S_0|}{|N|}\right) q + O(\sqrt{q})$$

where S_0 is the set of group elements in the coset xN which act as derangements on the set of roots of $f(T) - t = 0$. The constant in the above error term depends only on n , not on q .

As an example, let $f(T) = T^r$ with r prime and different from p (the characteristic of the base field \mathbb{F}_q). The Galois closure of $\mathbb{F}_q(T)/\mathbb{F}_q(T^r)$ is $\mathbb{F}_q(\mu, T)$ where μ is a nontrivial r th root of 1. Thus N is cyclic of order r and G/N is isomorphic to the Galois group of $\mathbb{F}_q(\mu)/\mathbb{F}_q$. The permutation action is of degree r . If $G = N$, then every non-trivial element is a derangement and so the image of f has order roughly $\frac{q}{r} + O(\sqrt{q})$. If $G \neq N$, then G is a Frobenius group and every fixed point free element is contained in N . Indeed, since in this case $(r, q-1) = 1$, we see that T^r is bijective on \mathbb{F}_q . For further examples, see Guralnick-Wan [GW] and references therein. Using work on derangements, they prove that if the degree of f is relatively prime to the characteristic p , then either f is bijective or $V_f \leq \frac{5q}{6} + O(\sqrt{q})$.

8.2. Proportion of derangements and Shalev's conjecture. Let G be a finite permutation group acting transitively on a set X of size $n > 1$. Subsection 8.1 motivated the study of $\delta(G, X)$, the proportion of derangements of G acting on X . We describe some results on this question, focusing particularly on lower bounds and analogs of our main results for classical groups.

Perhaps the earliest such result is due to Jordan [Jo], who showed that $\delta(G, X) > 0$. Cameron and Cohen [CaCo] proved that $\delta(G, X) \geq 1/n$ with equality if and only if G is a Frobenius group of order $n(n-1)$, where n is a prime power. See also [Se], who also notes a topological application of Jordan's theorem.

Based on extensive computations, it was asked in [BD] whether there is a universal constant $\delta > 0$ (which they speculate may be optimally chosen as $\frac{2}{7}$) such that $\delta(G, X) > \delta$ for all finite simple groups G . The existence of such a $\delta > 0$ has also been attributed as a conjecture of Shalev, who confirmed it for all actions of $PGL_d(q)$ with q fixed [Sh1].

Shalev's conjecture was proved by Fulman and Guralnick in the series of papers [FG1],[FG2],[FG3]. We do not attempt to sketch a proof of Shalev's conjecture here, but make a few remarks:

- (1) One can assume that the action of G on X is primitive, for if $f : Y \mapsto X$ is a surjection of G -sets, then $\delta(G, Y) \geq \delta(G, X)$.

- (2) By Jordan's theorem [Jo] that $\delta(G, X) > 0$, the proof of Shalev's conjecture is an asymptotic result: we only need to show that there exists a $\delta > 0$ such that for any sequence G_i, X_i with $|X_i| \rightarrow \infty$, one has that $\delta(G_i, X_i) > \delta$ for all sufficiently large i .
- (3) When G is the alternating group, by an argument similar to that of Theorem 3.3 of the current paper, the work of Luczak and Pyber implies that for all primitive actions of A_n except the action on k -sets, the proportion of derangements tends to 1. For the case of A_n on k -sets, one can give arguments similar to those Dixon [Dx1], who proved that the proportion of elements of S_n which are derangements on k -sets is at least $\frac{1}{3}$.
- (4) When G is a finite Chevalley group, the key is to study the set of regular semisimple elements of G . Typically (there are some exceptions in the orthogonal cases) this is the set of elements of G whose characteristic polynomial is square-free. Now a regular semisimple element is contained in a unique maximal torus, and there is a map from maximal tori to conjugacy classes of the Weyl group. This allows one to relate derangements in G to derangements in the Weyl group. For example, one concludes that the proportion of elements of $GL(n, q)$ which are regular semisimple and fix some k -space is at most the proportion of elements in S_n which fix a k -set. For large q , algebraic group arguments show that nearly all elements of $GL(n, q)$ are regular semisimple, and for fixed q , one uses generating functions to uniformly bound the proportion of regular semisimple elements away from 0.

To close this subsection, we note that the main result of this paper has an analog for finite classical groups. The following result was stated in [FG1] and is proved in [FG2].

Theorem 8.2. *Let G_i be a sequence of classical groups with the natural module of dimension d_i . Let X_i be a G_i -orbit of either totally singular or nondegenerate subspaces (of the natural module) of dimension $k_i \leq d_i/2$. If $k_i \rightarrow \infty$, then $\lim \delta(G_i, X_i) = 1$. If k_i is a bounded sequence, then there exist $0 < \delta_1 < \delta_2 < 1$ so that $\delta_1 < \delta(G_i, X_i) < \delta_2$.*

We remark that in characteristic 2, we consider the orthogonal group inside the symplectic group as the stabilizer of a subspace (indeed, if we view $Sp(2m, 2^e) = O(2m + 1, q^e)$, then the orthogonal groups are stabilizers of nondegenerate hyperplanes).

In fact, Fulman and Guralnick prove an analog of the Luczak-Pyber result for symmetric groups.

Theorem 8.3. *Let G_i be a sequence of classical groups with the natural module V_i of dimension d_i with $d_i \rightarrow \infty$. Let H_i be the union of all irreducible subgroups of G_i not containing G'_i (excluding orthogonal subgroups of the symplectic group in characteristic 2). Then $\lim_{i \rightarrow \infty} |H_i|/|G_i| = 0$.*

If the d_i are fixed, then this result is false. For example, if $G_i = PSL(2, q)$ and H is the normalizer of a maximal torus of G , then $\lim_{q \rightarrow \infty} \delta(G, G/H) = 1/2$. However, the analog of the previous theorem is proved in [FG1] if the rank of the Chevalley group is fixed. In this case, we take H_i to be the union of maximal subgroups which do not contain a maximal torus.

The example given above shows that the rank of the permutation action going to ∞ does not imply that the proportion of derangements tends to 1. The results of Fulman and Guralnick do show this is true if one considers simple Chevalley groups over fields of bounded size.

8.3. Fixed point ratios. Previous sections of this paper have discussed $fix(x)$, the number of fixed points of an element x of G on a set Ω . This subsection concerns the fixed point ratio $rfix(x) = \frac{fix(x)}{|\Omega|}$. We describe applications to random generation. For many other applications (base size, Guralnick-Thompson conjecture, etc.), see the survey [Sh2]. It should also be mentioned that fixed point ratios are a special case of character ratios, which have numerous applications to areas such as random walk [D] and number theory [GLM].

Let $P(G)$ denote the probability that two random elements of a finite group G generate G . One of the first results concerning $P(G)$ is due to Dixon [Dx2], who proved that $\lim_{n \rightarrow \infty} P(A_n) = 1$. The corresponding result for finite simple classical groups is due to Kantor and Lubotzky [KL]. The strategy adopted by Kantor and Lubotzky was to first note that for any pair $g, h \in G$, one has that $\langle g, h \rangle \neq G$ if and only if $\langle g, h \rangle$ is contained in a maximal subgroup M of G . Since $P(g, h \in M) = (|M|/|G|)^2$, it follows that

$$1 - P(G) \leq \sum_M \left(\frac{|M|}{|G|} \right)^2 \leq \sum_i \left(\frac{|M_i|}{|G|} \right)^2 \left(\frac{|G|}{|M_i|} \right) = \sum_i \frac{|M_i|}{|G|}.$$

Here M denotes a maximal subgroup and $\{M_i\}$ are representatives of conjugacy classes of maximal subgroups. Roughly, to show that this sum is small, one can use Aschbacher's classification of maximal subgroups [As], together with Liebeck's upper bounds on sizes of maximal subgroups [Li].

Now suppose that one wants to study $P_x(G)$, the chance that a fixed element x and a random element g of G generate G . Then

$$1 - P_x(G) = P(\langle x, g \rangle \neq G) \leq \sum_{\substack{M \text{ maximal} \\ x \in M}} P(g \in M) = \sum_{\substack{M \text{ maximal} \\ x \in M}} \frac{|M|}{|G|}.$$

Here the sum is over maximal subgroups M containing x . Let $\{M_i\}$ be a set of representatives of maximal subgroups of G , and write $M \sim M_i$ if M is conjugate to M_i . Then the above sum becomes

$$\sum_i \frac{|M_i|}{|G|} \sum_{\substack{M \sim M_i \\ x \in M}} 1.$$

To proceed further we assume that G is simple. Then, letting $N_G(M_i)$ denote the normalizer of M_i in G , one has that

$$g_1 M_i g_1^{-1} = g_2 M_i g_2^{-1} \leftrightarrow g_1^{-1} g_2 \in N_G(M_i) \leftrightarrow g_1^{-1} g_2 \in M_i \leftrightarrow g_1 M_i = g_2 M_i.$$

In other words, there is a bijection between conjugates of M_i and left cosets of M_i . Moreover, $x \in g M_i g^{-1}$ if and only if $x g M_i = g M_i$. Thus

$$\frac{|M_i|}{|G|} \sum_{\substack{M \sim M_i \\ x \in M}} 1 = \mathit{fix}(x, M_i).$$

Here $\mathit{fix}(x, M_i)$ denotes the fixed point ratio of x on left cosets of M_i , that is the proportion of left cosets of M_i fixed by x . Summarizing, $P_x(G)$ can be upper bounded in terms of the quantities $\mathit{fix}(x, M_i)$. This fact has been usefully applied in quite a few papers (see [GKa], [FG4] and the references therein, for example).

8.4. Miscellany. This subsection collects some miscellaneous facts about fixed points and derangements.

(1) *Formulae for fixed points*

We next state a well-known elementary proposition which gives different formulae for the number of fixed points of an element in a group action.

Proposition 8.4. *Let G be a finite group acting transitively on X . Let C be a conjugacy class of G and g in C . Let H be the stabilizer of a point in X .*

- (1) *The number of fixed points of g on X is $\frac{|C \cap H|}{|C|} |X|$.*
- (2) *The fixed point ratio of g on X is $\frac{|C \cap H|}{|C|}$.*
- (3) *The number of fixed points of g on X is $|C_G(g)| \sum_i |C_H(g_i)|^{-1}$ where the g_i are representatives for the H classes of $C \cap H$.*

Proof. Clearly (1) and (2) are equivalent. To prove (1), we determine the cardinality of the set $\{(u, x) \in C \times X \mid ux = x\}$. On the one hand, this set has size $|C|f(g)$ where $f(g)$ is the number of fixed points of g . On the other hand, it is $|X||C \cap H|$, whence (a) holds.

For (c), note that $|C| = \frac{|G|}{|C_G(g)|}$ and $|C \cap H| = \sum_i \frac{|H|}{|C_H(g_i)|}$ where the g_i are representative for the H -classes of $C \cap H$. Plugging this into (1) and using $|X| = \frac{|G|}{|H|}$ completes the proof. \square

(2) *Algorithmic issues*

It is natural to ask for an algorithm to generate a random derangement in S_n , for example for cryptographic purposes. Of course, one method is to simply generate random permutations until a derangement is reached. A more closed form algorithm has been suggested by Sam Payne. This begins by generating a random permutation and then, working left to right, each fixed point is transposed with a randomly chosen place. Each such transposition decreases the number of fixed points and an inductive argument shows

that after one pass, the resulting derangement is uniformly distributed. We do not know if this works starting with the identity permutation instead of a random permutation.

A very different, direct algorithm for generating a uniformly chosen derangement appears in [De]. There is also a literature on Gray codes for running through all derangements in the symmetric group; see [BV] and [KoL].

(3) *Algebraic combinatorics*

The set of derangements has itself been the subject of some combinatorial study. For example, Désarménien [De] has shown that there is a bijection between derangements in S_n and the set of permutations with first ascent occurring in an even position. This is extended and refined by Désarménien and Wachs [DeW]. Diaconis, McGrath, and Pitman [DMP] study the set of derangements with a single descent. They show that this set admits an associative, commutative product and unique factorization into cyclic elements. Bóna [Bn] studies the distribution of cycles in derangements, using among other things a result of E. Canfield that the associated generating function has all real zeros.

(4) *Statistics*

The fixed points of a permutation give rise to a useful metric on the permutation group: the Hamming metric. Thus $d(\pi, \sigma)$ is equal to the number of places where π and σ disagree. This is a bi-invariant metric on the permutation group and

$$d(\pi, \sigma) = d(id, \pi^{-1}\sigma) = \text{number of fixed points in } \pi^{-1}\sigma.$$

Such metrics have many statistical applications (Chapter 6 of [D]).

REFERENCES

- [Al] Aldous, D., *Probability approximations via the Poisson clumping heuristic*. Applied Mathematical Sciences, 77. Springer-Verlag, New York, 1989.
- [AT] Arratia, R., Tavaré, S., The cycle structure of random permutations, *Ann. Probab.* **20** (1992), 1567-1591.
- [As] Aschbacher, M., On the maximal subgroups of finite classical groups, *Invent. Math.* **76** (1984), 469-514.
- [AS] Aschbacher, M., Scott, L., Maximal subgroups of finite groups, *J. Algebra* **92** (1985), 44-80.
- [Ba] Babai, L., On the order of uniprimitive permutation groups, *Ann. of Math.* **113** (1981), 553-568.
- [BHJ] Barbour, A., Holst, L., Janson, S., *Poisson approximation*. Oxford Science Publications. Clarendon Press, Oxford University Press, New York, 1992.
- [BV] Baril, J., Vajnovszki, V., Gray code for derangements, *Discrete Appl. Math.* **140** (2004), 207-221.
- [BHR] Bidigare, P., Hanlon, P., Rockmore, D., A combinatorial description of the spectrum of the Tsetlin library and its generalization to hyperplane arrangements, *Duke Math. J.* **99** (1999), 135-174.

- [Bn] Bóna, M., On a balanced property of derangements, *Electron. J. Combin.* **13** (2006), Research Paper 102, 12 pp. (electronic).
- [BD] Boston, N., Dabrowski, T., Foguel, P., Gies, P., Leavitt, J., Ose, D., Jackson, D.A., The proportion of fixed-point-free elements of a transitive permutation group, *Comm. Algebra* **21** (1993), 3259-3275.
- [Bo] Bovey, J., The probability that some power of a permutation has small degree, *Bull. London Math. Soc.* **12** (1980), 47-51.
- [BLP] Buhler, J., Lenstra, H.W. Jr., Pomerance, C., Factoring integers with the number field sieve, in *The development of the number field sieve*, Lecture Notes in Mathematics **1554**, Springer-Verlag, Berlin, 1993.
- [Ca1] Cameron, P., *Permutation groups.*, London Mathematical Society Student Texts, 45. Cambridge University Press, 1999.
- [Ca2] Cameron, P., Derangements and p-elements in permutation groups, online lecture notes (2007), available at: <http://www.maths.qmul.ac.uk/~pjc/slides>.
- [CaCo] Cameron, P., Cohen, A.M., On the number of fixed point free elements in a permutation group, *Discrete Math.* **106/107** (1992), 135-138.
- [CDM] Chatterjee, S., Diaconis, P., Meckes, E., Exchangeable pairs and Poisson approximation, *Probab. Surv.* **2** (2005), 64-106 (electronic).
- [Ch] Chowla, S., The Riemann zeta and allied functions, *Bull. Amer. Math. Soc.* **58** (1952), 287-303.
- [Co] Cohen, S.D., The distribution of polynomials over finite fields, *Acta Arith.* **17** (1970), 255-271.
- [De] Désarménien, J., Une autre interprétation du nombre de dérangements, *Sém. Lothar. de Combin.* **1982**, B08b, 6 pp.
- [DeW] Désarménien, J., Wachs, M., Descent classes of permutations with a given number of fixed points, *J. Combin. Theory Ser. A* **64** (1993), 311-328.
- [D] Diaconis, P., *Group representations in probability and statistics*, Institute of Mathematical Statistics Lecture Notes 11, Hayward, CA, 1988.
- [DH1] Diaconis, P., Holmes, S., Matchings and phylogenetic trees, *Proc. Natl. Acad. Sci. USA* **95** (1998), 14600-14602.
- [DH2] Diaconis, P., Holmes, S., Random walk on trees and matchings, *Electronic Jour Probab.* **7** (2002), 1-17.
- [DMP] Diaconis, P., McGrath, M., Pitman, J., Riffle shuffles, cycles, and descents, *Combinatorica* **15** (1995), 11-29.
- [DS] Diaconis, P., Shahshahani, M., On the eigenvalues of random matrices, *J. Appl. Probab.* **31A** (1994), 49-62.
- [Dx1] Dixon, J., Random sets which invariably generate the symmetric group, *Discrete Math.* **105** (1992), 25-39.
- [Dx2] Dixon, J., The probability of generating the symmetric group, *Math. Z.* **110** (1969), 199-205.
- [DxM] Dixon, J., Mortimer, B., *Permutation groups*, Graduate Texts in Mathematics, 163. Springer-Verlag, 1996.
- [Du] Durrett, R., *Probability: theory and examples*, Second edition, Duxbury Press, 1996.
- [Fe] Feller, W., *An introduction to probability theory and its applications, Vol. 1*, Second edition, John Wiley & Sons, 1957.
- [F1] Fulman, J., Card shuffling and the decomposition of tensor products, *Pacific J. Math* **217** (2004), 247-262.
- [F2] Fulman, J., Convergence rates of random walk on irreducible representations of finite groups, *J. Theoret. Probab.*, to appear.
- [FG1] Fulman, J., Guralnick, R., Derangements in simple and primitive groups, in *Groups, combinatorics, and geometry: Durham 2001*, World Scientific Publishing, 2003, 99-121.

- [FG2] Fulman, J., Guralnick, R., Derangements in subspace actions of classical groups, preprint.
- [FG3] Fulman, J., Guralnick, R., Derangements in classical groups for non-subspace actions, preprint.
- [FG4] Fulman, J., Guralnick, R., The probability of generating an irreducible subgroup, preprint.
- [GIM] Gluck, D., Magaard, K., Absolute fourth moments and finiteness of linear groups, *Comm. Algebra* **34** (2006), 3387-3407.
- [Go] Goncharov, V., Sur la distribution des cycles dans les permutations, *C. R. (Doklady) Acad. Sci. URSS (N.S.)* **35** (1942), 267-269.
- [GKa] Guralnick, R., Kantor, W., Probabilistic generation of finite simple groups, *J. Algebra* **234** (2000), 743-792.
- [GKi] Guralnick, R., Kimmerle, W., On the cohomology of alternating and symmetric groups and decomposition of relation modules. *J. Pure Appl. Algebra* **69** (1990), 135-140.
- [GM] Guralnick, R., Magaard, K., On the minimal degree of a primitive permutation group, *J. Algebra* **207** (1998), 127-145.
- [GW] Guralnick, R., Wan, D., Bounds for fixed point free elements in a transitive group and applications to curves over finite fields, *Israel J. Math.* **101** (1997), 255-287.
- [JK] James, G., Kerber, A., *The representation theory of the symmetric group*, Encyclopedia of Mathematics and its Applications, 16, Addison-Wesley Publishing Co., Reading Mass., 1981.
- [Jo] Jordan, C., Recherches sur les substitutions, *J. Liouville* **17** (1872), 351-367.
- [KL] Kantor, W., Lubotzky, A., The probability of generating a finite classical group, *Geom. Dedicata* **36** (1990), 67-87.
- [KoL] Korsh, J., LaFollette, P., Constant time generation for derangements, *Inform. Process. Lett.* **90** (2004), 181-186.
- [KS] Kurzweil, H., Stellmacher, B., *The theory of finite groups: an introduction*, Springer-Verlag, 2004.
- [LS] Lenstra, H.W. Jr., Stevenhagen, P., Chebotarev and his density theorem, *Math. Intelligencer* **18** (1996), 26-37.
- [Li] Liebeck, M., On the order of maximal subgroups of finite classical groups, *Proc. London Math. Soc.* **50** (1985), 426-446.
- [LPS1] Liebeck, M., Praeger, C., Saxl, J., A classification of the maximal subgroups of the finite alternating and symmetric groups, *J. Algebra* **111** (1987), 365-383.
- [LPS2] Liebeck, M., Praeger, C., Saxl, J., On the O’Nan-Scott theorem for finite primitive permutation groups, *J. Austral. Math. Soc. Ser. A.* **44** (1988), 389-396.
- [LoPl] Lovász, L., Plummer, M., *Matching theory*, Annals of Discrete Mathematics 29, North Holland Publishing Co., Amsterdam, 1986.
- [LuPy] Luczak, T., Pyber, L., On random generation of the symmetric group, *Combin. Probab. Comput.* **2** (1993), 505-512.
- [Mo] Montmort, P.R. de, *Essay d’analyse sur les jeux de hazard*, (1708) 1st ed. , (1713) (2nd ed.). Jacques Quillau, Paris. Reprinted 1980 by Chelsea, New York.
- [Mor] Mortimer, B., Permutation groups containing affine groups of the same degree, *J. London Math. Soc.* **15** (1977), 445-455.
- [NB] Newton, B., Benesh, B., A classification of certain maximal subgroups of symmetric groups, *J. Algebra* **304** (2006), 1108-1113.
- [O] Odlyzko, A.M., Asymptotic enumeration methods, Chapter 22 of Handbook of Combinatorics, Volume II. MIT Press and Elsevier, 1995.
- [P] Passman, D., *Permutation Groups*, W. A. Benjamin, Inc., New York-Amsterdam 1968.
- [Se] Serre, J.P., On a theorem of Jordan, *Bull. Amer. Math. Soc.* **40** (2003), 429-440.

- [Sh1] Shalev, A., A theorem on random matrices and some applications, *J. Algebra* **199**, 124-141.
- [Sh2] Shalev, A., Probabilistic group theory, in *Groups St. Andrews 1997*, London Math. Soc. Lecture Note Ser. 261, Cambridge Univ. Press., Cambridge, 1999, 648-678.
- [SL] Shepp, L., Lloyd. S., Ordered cycle lengths in a random permutation, *Trans. Amer. Math. Soc.* **121** (1966), 340-357.
- [St] Stanley, R., Enumerative combinatorics. Vol. 1. Corrected reprint of the 1986 original, Cambridge Studies in Advanced Mathematics, 49, Cambridge University Press, Cambridge, 1997.
- [Ta] Takács, L., The problem of coincidences, *Arch. Hist. Exact Sci.* **21** (1979/80), no. 3, 229-244.

DEPARTMENT OF MATHEMATICS AND STATISTICS, STANFORD UNIVERSITY, STANFORD, CA 94305

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SOUTHERN CALIFORNIA, LOS ANGELES, CA 90089-2532

E-mail address: `fulman@usc.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SOUTHERN CALIFORNIA, LOS ANGELES, CA 90089-2532

E-mail address: `guralnic@usc.edu`