

Lecture 15: Logistic and Poisson Regression

Nancy R. Zhang

Statistics 191, Stanford University

March 10, 2008

Review - Binary responses model

Model: $Y \in \{0, 1\}$,

$$P(Y = 1 | X_1, \dots, X_p) = g^{-1}(\beta_1 X_1 + \beta_2 X_2 + \dots + \beta_p X_p).$$

Where

$$g(\pi) = \log\left(\frac{\pi}{1 - \pi}\right).$$

The inverse g^{-1} is

$$g^{-1}(z) = \frac{e^z}{1 + e^z}.$$

We have no choice but to accept non-constant variance,

$$\text{Var}(Y) = \pi(X)[1 - \pi(X)].$$

Review - Model interpretation

An intuitive quantity to assess probabilities:

$$odds = \frac{P(Y = 1|X)}{P(Y = 0|X)}.$$

In the logistic regression model,

$$\log(odds) = \beta X.$$

The parameter β is the contribution of unit increase in X to the increase (decrease) in odds. For example, if X were binary as well,

$$\log\left(\frac{odds(X = 1)}{odds(X = 0)}\right) = \beta.$$

Logit Model for Multinomial Response

If the response Y belong to K categories.

① Designate one category as the “base” category.

②

$$P(Y = k|X) = \frac{e^{X\beta_k}}{1 + \sum_{l=1}^{K-1} e^{X\beta_l}}$$

Here, $\beta_k = (\beta_{k1}, \dots, \beta_{kp})$.

$$P(Y = K|X) = \frac{1}{1 + \sum_{l=1}^{K-1} e^{X\beta_l}}$$

③ $p \times (K - 1)$ parameters.

④ β_{ki} for k -th category and i -th predictor interpreted as increase in log-odds from base category.

Logit Model for Multinomial Response

Equivalent definition:

$$\log \frac{\pi_k(X)}{\pi_K(X)} = \alpha_k + X\beta_k, \quad k = 1, \dots, K - 1,$$

where

$$\pi_k(X) = P(Y = k|X).$$

Alligator Food Example

Study on the primary food choice of alligators.

- 1 Data: 219 alligators captured in four Florida lakes.
- 2 Response variable: food type, in volume, found in the alligator's stomach. 5 categories:
 - 1 fish
 - 2 invertebrate
 - 3 reptile
 - 4 bird
 - 5 other
- 3 Predictors:
 - 1 Lake of capture (Hancock, Oklawaha, Trafford, George)
 - 2 Gender (M, F).
 - 3 Size ($\leq 2.3m$, $\geq 2.3m$).

Alligator Food Choice Example

TABLE 7.1 Primary Food Choice of Alligators

Lake	Gender	Size (m)	Primary Food Choice				
			Fish	Invertebrate	Reptile	Bird	Other
Hancock	Male	≤ 2.3	7	1	0	0	5
		> 2.3	4	0	0	1	2
	Female	≤ 2.3	16	3	2	2	3
		> 2.3	3	0	1	2	3
Oklawaha	Male	≤ 2.3	2	2	0	0	1
		> 2.3	13	7	6	0	0
	Female	≤ 2.3	3	9	1	0	2
		> 2.3	0	1	0	1	0
Trafford	Male	≤ 2.3	3	7	1	0	1
		> 2.3	8	6	6	3	5
	Female	≤ 2.3	2	4	1	1	4
		> 2.3	0	1	0	0	0
George	Male	≤ 2.3	13	10	0	2	2
		> 2.3	9	0	0	1	2
	Female	≤ 2.3	3	9	1	0	1
		> 2.3	8	1	0	0	1

Source: Data courtesy of Clint Moore, from an unpublished manuscript by M. F. Delaney and C. T. Moore.

- 1 Do gender, size, or lake of capture influence food choice?
- 2 Are there interaction effects?
- 3 Obtain estimates of $P(\text{food choice} = \text{fish} \mid \text{Gender, Size, Lake})$.

Functions for multinomial fitting in R: `multinom` in library `nnet`.

Fitted probabilities \hat{p}_i

If you had data about the size of the alligators (and not just the classification (\leq or ≥ 2.3 m)), then you can estimate a response curve like this:




FIGURE 7.1 Estimated probabilities for primary food choice.

From Agresti, Categorical Data Analysis

Review: 2×2 tables: Gender and After-life

	Y	N or U	
M	435	147	582
F	375	134	509
Total	810	281	1091

- 1 Poisson sampling assumption: row sums not fixed.

Review: 2×2 tables: Gender and After-life

	Y	N or U	
M	435	147	582
F	375	134	509
Total	810	281	1091

- 1 Poisson sampling assumption: row sums not fixed.
- 2 Model: $Y_{i,j} \sim \text{Poisson}(\lambda_{ij})$.

Review: 2×2 tables: Gender and After-life

	Y	N or U	
M	435	147	582
F	375	134	509
Total	810	281	1091

- 1 Poisson sampling assumption: row sums not fixed.
- 2 Model: $Y_{i,j} \sim \text{Poisson}(\lambda_{ij})$.
- 3 H_0 : Row and columns independent (i.e. $\log \lambda_{ij} = \lambda + \alpha_i + \beta_j$).
(Before I used $\lambda_{ij} = \lambda \alpha_i \beta_j$, now I get rid of log to save space).

Review: 2×2 tables: Gender and After-life

	Y	N or U	
M	435	147	582
F	375	134	509
Total	810	281	1091

- 1 Poisson sampling assumption: row sums not fixed.
- 2 Model: $Y_{i,j} \sim \text{Poisson}(\lambda_{ij})$.
- 3 H_0 : Row and columns independent (i.e. $\log \lambda_{ij} = \lambda + \alpha_i + \beta_j$).
(Before I used $\lambda_{ij} = \lambda \alpha_i \beta_j$, now I get rid of log to save space).

This is the simplest type of contingency table.

3-way tables: Alcohol Cigarette, and Marijuana Use

Survey asked 2276 students in their final year of high school in a nonurban area near Dayton, Ohio whether they ever used alcohol, cigarettes, or marijuana.

Alcohol Use	Cigarette Use	Marijuana Use	
		Yes	No
Yes	Yes	911	538
	No	44	456
No	Yes	3	43
	No	2	279

This is example of a $2 \times 2 \times 2$ contingency table. Shorthand: A=alcohol, C=cigarette, M=marijuana.

3-way tables: Types of Interaction

Alcohol Use	Cigarette Use	Marijuana Use	
		Yes	No
Yes	Yes	911	538
	No	44	456
No	Yes	3	43
	No	2	279

3-way tables: Types of Interaction

Alcohol Use	Cigarette Use	Marijuana Use	
		Yes	No
Yes	Yes	911	538
	No	44	456
No	Yes	3	43
	No	2	279

$$Y_{ijk} \sim \text{Poisson}(\lambda_{ijk})$$

3-way tables: Types of Interaction

Alcohol Use	Cigarette Use	Marijuana Use	
		Yes	No
Yes	Yes	911	538
	No	44	456
No	Yes	3	43
	No	2	279

$$Y_{ijk} \sim \text{Poisson}(\lambda_{ijk})$$

Conditioned on total (N) $Y_{ijk} \sim \text{Multinom}(N, \pi_{ijk})$.

3-way tables: Types of Interaction

Alcohol Use	Cigarette Use	Marijuana Use	
		Yes	No
Yes	Yes	911	538
	No	44	456
No	Yes	3	43
	No	2	279

$$Y_{ijk} \sim \text{Poisson}(\lambda_{ijk})$$

Conditioned on total (N) $Y_{ijk} \sim \text{Multinom}(N, \pi_{ijk})$.

π_{i++} be probability of row $A = i$,

π_{ij+} be probability of $A = i, C = j$, etc.

3-way tables: Types of Interaction

Alcohol Use	Cigarette Use	Marijuana Use	
		Yes	No
Yes	Yes	911	538
	No	44	456
No	Yes	3	43
	No	2	279

$$Y_{ijk} \sim \text{Poisson}(\lambda_{ijk})$$

Conditioned on total (N) $Y_{ijk} \sim \text{Multinom}(N, \pi_{ijk})$.

π_{i++} be probability of row $A = i$,

π_{ij+} be probability of $A = i, C = j$, etc.

1 A,C, and M mutually independent

$$\log \lambda_{ijk} = \lambda + \lambda_i^A + \lambda_j^C + \lambda_k^M$$

$$\pi_{ijk} = \pi_{i++}\pi_{+j+}\pi_{++k}$$

3-way tables: Types of Interaction

Alcohol Use	Cigarette Use	Marijuana Use	
		Yes	No
Yes	Yes	911	538
	No	44	456
No	Yes	3	43
	No	2	279

$$Y_{ijk} \sim \text{Poisson}(\lambda_{ijk})$$

Conditioned on total (N) $Y_{ijk} \sim \text{Multinom}(N, \pi_{ijk})$.

π_{i++} be probability of row $A = i$,

π_{ij+} be probability of $A = i, C = j$, etc.

- 1 M is **jointly independent** of A, C

$$\log \lambda_{ijk} = \lambda + \lambda_i^A + \lambda_j^C + \lambda_k^M + \lambda_{ij}^{AC}$$

$$\pi_{ijk} = \pi_{ij+} \pi_{++k}$$

3-way tables: Types of Interaction

Alcohol Use	Cigarette Use	Marijuana Use	
		Yes	No
Yes	Yes	911	538
	No	44	456
No	Yes	3	43
	No	2	279

$$Y_{ijk} \sim \text{Poisson}(\lambda_{ijk})$$

Conditioned on total (N) $Y_{ijk} \sim \text{Multinom}(N, \pi_{ijk})$.

π_{i++} be probability of row $A = i$,

π_{ij+} be probability of $A = i, C = j$, etc.

1 C and M **conditionally independent** given A

$$\log \lambda_{ijk} = \lambda + \lambda_i^A + \lambda_j^C + \lambda_k^M + \lambda_{ij}^{AC} + \lambda_{ik}^{AM}$$

$$\pi_{jk|i} = \pi_{j+|i} \pi_{+k|i}$$

3-way tables: Types of Interaction

Alcohol Use	Cigarette Use	Marijuana Use	
		Yes	No
Yes	Yes	911	538
	No	44	456
No	Yes	3	43
	No	2	279

$$Y_{ijk} \sim \text{Poisson}(\lambda_{ijk})$$

Conditioned on total (N) $Y_{ijk} \sim \text{Multinom}(N, \pi_{ijk})$.

π_{i++} be probability of row $A = i$,

π_{ij+} be probability of $A = i, C = j$, etc.

- 1 Each pair of A,C, and M has **homogeneous association**.

$$\log \lambda_{ijk} = \lambda + \lambda_i^A + \lambda_j^C + \lambda_k^M + \lambda_{ij}^{AC} + \lambda_{ik}^{AM} + \lambda_{ik}^{CM}.$$

e.g. the dependence relationship of A, C does not depend on M.

3-way tables: Types of Interaction

Alcohol Use	Cigarette Use	Marijuana Use	
		Yes	No
Yes	Yes	911	538
	No	44	456
No	Yes	3	43
	No	2	279

$$Y_{ijk} \sim \text{Poisson}(\lambda_{ijk})$$

Conditioned on total (N) $Y_{ijk} \sim \text{Multinom}(N, \pi_{ijk})$.

π_{i++} be probability of row $A = i$,


π_{ij+} be probability of $A = i, C = j$, etc.

1 Saturated Model.

$$\log \lambda_{ijk} = \lambda + \lambda_i^A + \lambda_j^C + \lambda_k^M + \lambda_{ij}^{AC} + \lambda_{ik}^{AM} + \lambda_{jk}^{CM} + \lambda_{ijk}^{ACM}.$$

3-way tables: Types of Interaction

Symbol	Interpretation
(A,C,M)	Mutual Independence
(AC,M)	AC jointly independent of M
(AC,AM)	M, C conditionally independent given A
(AC,AM,CM)	Homogeneous association of each pair.



Marginal independence: fit 2×2 table.

Analysis of 3-way tables

- 1 Fit log-linear model (Poisson GLM) for each of the models.
 - 1 Criterion: maximum likelihood.

Analysis of 3-way tables

- 1 Fit log-linear model (Poisson GLM) for each of the models.
 - 1 Criterion: maximum likelihood.
 - 2 Fitting method: Newton Raphson.

Analysis of 3-way tables

- 1 Fit log-linear model (Poisson GLM) for each of the models.
 - 1 Criterion: maximum likelihood.
 - 2 Fitting method: Newton Raphson.
- 2 Use a model selection criterion to choose the best one.
 - 1 AIC, BIC.

Analysis of 3-way tables

- 1 Fit log-linear model (Poisson GLM) for each of the models.
 - 1 Criterion: maximum likelihood.
 - 2 Fitting method: Newton Raphson.
- 2 Use a model selection criterion to choose the best one.
 - 1 AIC, BIC.
 - 2 Use Deviance χ^2 test to choose between nested models.

Analysis of 3-way tables

- 1 Fit log-linear model (Poisson GLM) for each of the models.
 - 1 Criterion: maximum likelihood.
 - 2 Fitting method: Newton Raphson.
- 2 Use a model selection criterion to choose the best one.
 - 1 AIC, BIC.
 - 2 Use Deviance χ^2 test to choose between nested models.

`glm(..., family=poisson), loglm(MASS)` in R.